

In the claims:

Please amend the following claims as indicated.

1-27. (Cancelled)

28. (Currently amended) A method for inducing an immune response ~~a host to produce Ab3 antibodies that specifically bind to prostate specific antigen~~ comprising administering to the ~~host an Ab1~~ a mammal an antibody or antigen-binding fragment thereof that specifically binds to an epitope of circulating prostate specific antigen, said epitope being an epitope to which ~~wherein the epitope is specifically bound by a monoclonal antibody produced by a hybridoma that has ATCC Designation Number HB-12526~~ specifically binds.

29. (Cancelled)

30. (Currently amended) The method of claim 28, wherein the ~~Ab1~~ antibody is a monoclonal antibody ~~produced by a hybridoma that has ATCC Designation Number HB-12526~~.

31. (Cancelled)

32. (Currently amended) The method of claim 28, wherein the ~~Ab1~~ antibody or antigen-binding fragment thereof is conjugated to an immunogenic carrier.

33. (Previously presented) The method of claim 32, wherein the immunogenic carrier is keyhole limpet hemocyanin.

34-35. (Cancelled)

36. (Currently amended) The method of claim 28, wherein the ~~Ab1~~ antibody is selected from the group consisting of ~~one member of an immunologic pair; an antibody; a monoclonal antibody, [[;]] an antibody fragment thereof; a single chain antibody [[;]], a humanized antibody, or fragment thereof; and a chimeric antibody or fragment thereof.~~

37. **(Currently amended)** The method according to claim ~~17~~28, wherein the ~~Ab1~~  
antibody or antigen binding fragment thereof is a xenogenic antibody or antigen-binding  
fragment thereof.
38. **(Currently amended)** The method according to claim ~~17~~28, wherein the ~~Ab1~~  
antibody or antigen-binding fragment thereof is formulated with an adjuvant.
39. **(Cancelled)**
40. **(Currently amended)** The method of claim ~~39~~28, wherein the ~~low-dose~~antibody or  
antigen-binding fragment thereof is formulated at a dose of from about 0.1 µg to about 2 mg per  
kilogram of body weight of the ~~patient~~mammal.
41. **(Currently amended)** The method of claim ~~39~~28, wherein the ~~low-dose~~antibody or  
antigen-binding fragment thereof is formulated at a dose of from about 1.0 µg to about 200 µg  
per kilogram of body weight of the ~~patient~~mammal.